

Search for Charge-1/3 Third-Generation Leptoquarks in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV

B. Abbott,³¹ M. Abolins,²⁷ B. S. Acharya,⁴⁶ I. Adam,¹² D. L. Adams,⁴⁰ M. Adams,¹⁷ S. Ahn,¹⁴ H. Aihara,²³ G. A. Alves,¹⁰ N. Amos,²⁶ E. W. Anderson,¹⁹ R. Astur,⁴⁵ M. M. Baarmand,⁴⁵ L. Babukhadia,² A. Baden,²⁵ V. Balamurali,³⁵ J. Balderston,¹⁶ B. Baldin,¹⁴ S. Banerjee,⁴⁶ J. Bantly,⁵ E. Barberis,²³ J. F. Bartlett,¹⁴ A. Belyaev,²⁹ S. B. Beri,³⁷ I. Bertram,³⁴ V. A. Bezzubov,³⁸ P. C. Bhat,¹⁴ V. Bhatnagar,³⁷ M. Bhattacharjee,⁴⁵ N. Biswas,³⁵ G. Blazey,³³ S. Blessing,¹⁵ P. Bloom,⁷ A. Boehnlein,¹⁴ N. I. Bojko,³⁸ F. Borchering,¹⁴ C. Boswell,⁹ A. Brandt,¹⁴ R. Brock,²⁷ A. Bross,¹⁴ D. Buchholz,³⁴ V. S. Burtovoi,³⁸ J. M. Butler,³ W. Carvalho,¹⁰ D. Casey,²⁷ Z. Casilum,⁴⁵ H. Castilla-Valdez,¹¹ D. Chakraborty,⁴⁵ S.-M. Chang,³² S. V. Chekulaev,³⁸ L.-P. Chen,²³ W. Chen,⁴⁵ S. Choi,⁴⁴ S. Chopra,²⁶ B. C. Choudhary,⁹ J. H. Christenson,¹⁴ M. Chung,¹⁷ D. Claes,³⁰ A. R. Clark,²³ W. G. Cobau,²⁵ J. Cochran,⁹ L. Coney,³⁵ W. E. Cooper,¹⁴ C. Cretsinger,⁴² D. Cullen-Vidal,⁵ M. A. C. Cummings,³³ D. Cutts,⁵ O. I. Dahl,²³ K. Davis,² K. De,⁴⁷ K. Del Signore,²⁶ M. Demarteau,¹⁴ D. Denisov,¹⁴ S. P. Denisov,³⁸ H. T. Diehl,¹⁴ M. Diesburg,¹⁴ G. Di Loreto,²⁷ P. Draper,⁴⁷ Y. Ducros,⁴³ L. V. Dudko,²⁹ S. R. Dugad,⁴⁶ D. Edmunds,²⁷ J. Ellison,⁹ V. D. Elvira,⁴⁵ R. Engelmann,⁴⁵ S. Eno,²⁵ G. Eppley,⁴⁰ P. Ermolov,²⁹ O. V. Eroshin,³⁸ V. N. Evdokimov,³⁸ T. Fahland,⁸ M. K. Fatyga,⁴² S. Feher,¹⁴ D. Fein,² T. Ferbel,⁴² G. Finocchiaro,⁴⁵ H. E. Fisk,¹⁴ Y. Fisyak,⁴ E. Flattum,¹⁴ G. E. Forden,² M. Fortner,³³ K. C. Frame,²⁷ S. Fuess,¹⁴ E. Gallas,⁴⁷ A. N. Galyaev,³⁸ P. Gartung,⁹ V. Gavrilov,²⁸ T. L. Geld,²⁷ R. J. Genik II,²⁷ K. Genser,¹⁴ C. E. Gerber,¹⁴ Y. Gershtein,²⁸ B. Gibbard,⁴ S. Glenn,⁷ B. Gobbi,³⁴ A. Goldschmidt,²³ B. Gómez,¹ G. Gómez,²⁵ P. I. Goncharov,³⁸ J. L. González Solís,¹¹ H. Gordon,⁴ L. T. Goss,⁴⁸ K. Gounder,⁹ A. Goussiou,⁴⁵ N. Graf,⁴ P. D. Grannis,⁴⁵ D. R. Green,¹⁴ H. Greenlee,¹⁴ S. Grinstein,⁶ P. Grudberg,²³ S. Grünendahl,¹⁴ G. Guglielmo,³⁶ J. A. Guida,² J. M. Guida,⁵ A. Gupta,⁴⁶ S. N. Gurzhiev,³⁸ G. Gutierrez,¹⁴ P. Gutierrez,³⁶ N. J. Hadley,²⁵ H. Haggerty,¹⁴ S. Hagopian,¹⁵ V. Hagopian,¹⁵ K. S. Hahn,⁴² R. E. Hall,⁸ P. Hanlet,³² S. Hansen,¹⁴ J. M. Hauptman,¹⁹ D. Hedin,³³ A. P. Heinson,⁹ U. Heintz,¹⁴ R. Hernández-Montoya,¹¹ T. Heuring,¹⁵ R. Hirosky,¹⁷ J. D. Hobbs,⁴⁵ B. Hoeneisen,^{1,*} J. S. Hoftun,⁵ F. Hsieh,²⁶ Ting Hu,⁴⁵ Tong Hu,¹⁸ T. Huehn,⁹ A. S. Ito,¹⁴ E. James,² J. Jaques,³⁵ S. A. Jerger,²⁷ R. Jesik,¹⁸ J. Z.-Y. Jiang,⁴⁵ T. Joffe-Minor,³⁴ K. Johns,² M. Johnson,¹⁴ A. Jonckheere,¹⁴ M. Jones,¹⁶ H. Jöstlein,¹⁴ S. Y. Jun,³⁴ C. K. Jung,⁴⁵ S. Kahn,⁴ G. Kalbfleisch,³⁶ J. S. Kang,²⁰ D. Karmanov,²⁹ D. Karmgard,¹⁵ R. Kehoe,³⁵ M. L. Kelly,³⁵ C. L. Kim,²⁰ S. K. Kim,⁴⁴ B. Klima,¹⁴ C. Klopfenstein,⁷ J. M. Kohli,³⁷ D. Koltick,³⁹ A. V. Kostritskiy,³⁸ J. Kotcher,⁴ A. V. Kotwal,¹² J. Kourlas,³¹ A. V. Kozelov,³⁸ E. A. Kozlovsky,³⁸ J. Krane,³⁰ M. R. Krishnaswamy,⁴⁶ S. Krzywdzinski,¹⁴ S. Kuleshov,²⁸ S. Kunori,²⁵ F. Landry,²⁷ G. Landsberg,¹⁴ B. Lauer,¹⁹ A. Leflat,²⁹ H. Li,⁴⁵ J. Li,⁴⁷ Q. Z. Li-Demarteau,¹⁴ J. G. R. Lima,⁴¹ D. Lincoln,¹⁴ S. L. Linn,¹⁵ J. Linnemann,²⁷ R. Lipton,¹⁴ Y. C. Liu,³⁴ F. Lobkowicz,⁴² S. C. Loken,²³ S. Lökös,⁴⁵ L. Lueking,¹⁴ A. L. Lyon,²⁵ A. K. A. Maciel,¹⁰ R. J. Madras,²³ R. Madden,¹⁵ L. Magaña-Mendoza,¹¹ V. Manankov,²⁹ S. Mani,⁷ H. S. Mao,^{14,†} R. Markeloff,³³ T. Marshall,¹⁸ M. I. Martin,¹⁴ K. M. Mauritz,¹⁹ B. May,³⁴ A. A. Mayorov,³⁸ R. McCarthy,⁴⁵ J. McDonald,¹⁵ T. McKibben,¹⁷ J. McKinley,²⁷ T. McMahon,³⁶ H. L. Melanson,¹⁴ M. Merkin,²⁹ K. W. Merritt,¹⁴ H. Miettinen,⁴⁰ A. Mincer,³¹ C. S. Mishra,¹⁴ N. Mokhov,¹⁴ N. K. Mondal,⁴⁶ H. E. Montgomery,¹⁴ P. Mooney,¹ H. da Motta,¹⁰ C. Murphy,¹⁷ F. Nang,² M. Narain,¹⁴ V. S. Narasimham,⁴⁶ A. Narayanan,² H. A. Neal,²⁶ J. P. Negret,¹ P. Nemethy,³¹ D. Norman,⁴⁸ L. Oesch,²⁶ V. Oguri,⁴¹ E. Oliveira,¹⁰ E. Oltman,²³ N. Oshima,¹⁴ D. Owen,²⁷ P. Padley,⁴⁰ A. Para,¹⁴ Y. M. Park,²¹ R. Partridge,⁵ N. Parua,⁴⁶ M. Paterno,⁴² B. Pawlik,²² J. Perkins,⁴⁷ M. Peters,¹⁶ R. Piegaia,⁶ H. Piekarczyk,¹⁵ Y. Pischalnikov,³⁹ B. G. Pope,²⁷ H. B. Prosper,¹⁵ S. Protopopescu,⁴ J. Qian,²⁶ P. Z. Quintas,¹⁴ R. Raja,¹⁴ S. Rajagopalan,⁴ O. Ramirez,¹⁷ L. Rasmussen,⁴⁵ S. Reucroft,³² M. Rijssenbeek,⁴⁵ T. Rockwell,²⁷ M. Roco,¹⁴ P. Rubinov,³⁴ R. Ruchti,³⁵ J. Rutherford,² A. Sánchez-Hernández,¹¹ A. Santoro,¹⁰ L. Sawyer,²⁴ R. D. Schamberger,⁴⁵ H. Schellman,³⁴ J. Sculli,³¹ E. Shabalina,²⁹ C. Shaffer,¹⁵ H. C. Shankar,⁴⁶ R. K. Shivpuri,¹³ M. Shupe,² H. Singh,⁹ J. B. Singh,³⁷ V. Sirotenko,³³ W. Smart,¹⁴ E. Smith,³⁶ R. P. Smith,¹⁴ R. Snihur,³⁴ G. R. Snow,³⁰ J. Snow,³⁶ S. Snyder,⁴ J. Solomon,¹⁷ M. Sosebee,⁴⁷ N. Sotnikova,²⁹ M. Souza,¹⁰ A. L. Spadafora,²³ G. Steinbrück,³⁶ R. W. Stephens,⁴⁷ M. L. Stevenson,²³ D. Stewart,²⁶ F. Stichelbaut,⁴⁵ D. Stoker,⁸ V. Stolin,²⁸ D. A. Stoyanova,³⁸ M. Strauss,³⁶ K. Streets,³¹ M. Strovink,²³ A. Sznajder,¹⁰ P. Tamburello,²⁵ J. Tarazi,⁸ M. Tartaglia,¹⁴ T. L. T. Thomas,³⁴ J. Thompson,²⁵ T. G. Trippe,²³ P. M. Tuts,¹² N. Varelas,¹⁷ E. W. Varnes,²³ D. Vititoe,² A. A. Volkov,³⁸ A. P. Vorobiev,³⁸ H. D. Wahl,¹⁵ G. Wang,¹⁵ J. Warchol,³⁵ G. Watts,⁵ M. Wayne,³⁵ H. Weerts,²⁷ A. White,⁴⁷ J. T. White,⁴⁸ J. A. Wightman,¹⁹ S. Willis,³³ S. J. Wimpenny,⁹ J. V. D. Wirjawan,⁴⁸ J. Womersley,¹⁴ E. Won,⁴² D. R. Wood,³² H. Xu,⁵ R. Yamada,¹⁴ P. Yamin,⁴ J. Yang,³¹ T. Yasuda,³² P. Yepes,⁴⁰ C. Yoshikawa,¹⁶ S. Youssef,¹⁵ J. Yu,¹⁴ Y. Yu,⁴⁴ Z. Zhou,¹⁹ Z. H. Zhu,⁴² D. Zieminska,¹⁸ A. Zieminski,¹⁸ E. G. Zverev,²⁹ and A. Zylberstein⁴³

(D0 Collaboration)

- ¹Universidad de los Andes, Bogotá, Colombia
²University of Arizona, Tucson, Arizona 85721
³Boston University, Boston, Massachusetts 02215
⁴Brookhaven National Laboratory, Upton, New York 11973
⁵Brown University, Providence, Rhode Island 02912
⁶Universidad de Buenos Aires, Buenos Aires, Argentina
⁷University of California, Davis, California 95616
⁸University of California, Irvine, California 92697
⁹University of California, Riverside, California 92521
¹⁰LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil
¹¹CINVESTAV, Mexico City, Mexico
¹²Columbia University, New York, New York 10027
¹³Delhi University, Delhi, India 110007
¹⁴Fermi National Accelerator Laboratory, Batavia, Illinois 60510
¹⁵Florida State University, Tallahassee, Florida 32306
¹⁶University of Hawaii, Honolulu, Hawaii 96822
¹⁷University of Illinois at Chicago, Chicago, Illinois 60607
¹⁸Indiana University, Bloomington, Indiana 47405
¹⁹Iowa State University, Ames, Iowa 50011
²⁰Korea University, Seoul, Korea
²¹Kyungshung University, Pusan, Korea
²²Institute of Nuclear Physics, Kraków, Poland
²³Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720
²⁴Louisiana Tech University, Ruston, Louisiana 71272
²⁵University of Maryland, College Park, Maryland 20742
²⁶University of Michigan, Ann Arbor, Michigan 48109
²⁷Michigan State University, East Lansing, Michigan 48824
²⁸Institute for Theoretical and Experimental Physics, Moscow, Russia
²⁹Moscow State University, Moscow, Russia
³⁰University of Nebraska, Lincoln, Nebraska 68588
³¹New York University, New York, New York 10003
³²Northeastern University, Boston, Massachusetts 02115
³³Northern Illinois University, DeKalb, Illinois 60115
³⁴Northwestern University, Evanston, Illinois 60208
³⁵University of Notre Dame, Notre Dame, Indiana 46556
³⁶University of Oklahoma, Norman, Oklahoma 73019
³⁷University of Panjab, Chandigarh 16-00-14, India
³⁸Institute for High Energy Physics, Protvino 142284, Russia
³⁹Purdue University, West Lafayette, Indiana 47907
⁴⁰Rice University, Houston, Texas 77005
⁴¹Universidade do Estado do Rio de Janeiro, Brazil
⁴²University of Rochester, Rochester, New York 14627
⁴³CEA, DAPNIA/Service de Physique des Particules, CE-Saclay, Gif-sur-Yvette, France
⁴⁴Seoul National University, Seoul, Korea
⁴⁵State University of New York, Stony Brook, New York 11794
⁴⁶Tata Institute of Fundamental Research, Colaba, Mumbai 400005, India
⁴⁷University of Texas, Arlington, Texas 76019
⁴⁸Texas A&M University, College Station, Texas 77843

(Received 10 March 1998)

We report on a search for charge-1/3 third-generation leptoquarks (LQ) produced in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV using the D0 detector at Fermilab. Third-generation leptoquarks are assumed to be produced in pairs and to decay to a tau neutrino and a b quark with branching fraction B . We place upper limits on $\sigma(p\bar{p} \rightarrow LQ\bar{L}Q)B^2$ as a function of the leptoquark mass M_{LQ} . Assuming $B = 1$, we exclude at the 95% confidence level third-generation scalar leptoquarks with $M_{LQ} < 94$ GeV/ c^2 , and third-generation vector leptoquarks with $M_{LQ} < 216$ GeV/ c^2 ($M_{LQ} < 148$ GeV/ c^2) assuming Yang-Mills (anomalous) coupling. [S0031-9007(98)06513-2]

PACS numbers: 14.80.-j, 13.85.Rm

Leptoquarks (LQ) are bosons predicted in many extensions to the standard model [1]. They carry both lepton and color quantum numbers, couple to leptons and quarks, and decay via $LQ \rightarrow l + q$. To satisfy experimental constraints on flavor changing neutral currents, leptoquarks of mass accessible to current collider experiments are constrained to couple to only one generation of leptons and quarks [2]. Therefore, only leptoquarks which couple within a single generation are considered here.

This Letter reports the results of a search for charge-1/3 third-generation leptoquarks produced in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. We assume that leptoquarks are produced in pairs by QCD processes such as $p\bar{p} \rightarrow g \rightarrow LQ\bar{L}Q + X$. This process dominates over other production mechanisms which depend on the unknown leptoquark-lepton-quark coupling λ under the standard condition $\lambda \leq \sqrt{4\pi}\alpha_{EM}$. We search for the decay signature where both leptoquarks decay via $LQ \rightarrow \nu_\tau + b$ resulting in a $\nu_\tau\bar{\nu}_\tau b\bar{b}$ final state. For a leptoquark mass (M_{LQ}) smaller than the mass of the top quark (m_t), the decay $LQ \rightarrow \tau + t$ is forbidden, and the branching fraction for the $\nu_\tau b$ mode, B , is unity [3]. For $M_{LQ} > m_t$, phase space factors suppress τt decays relative to the $\nu_\tau b$ channel. In this paper we give limits on the pair production cross section times B^2 (σB^2) for M_{LQ} between 50 and 300 GeV/ c^2 . Limits on the cross section are used to set limits on the third-generation leptoquark mass for scalar and vector leptoquarks. Previous limits from the LEP e^+c^- collider exclude all third-generation leptoquarks with masses below 45 GeV/ c^2 [4], while the CDF Collaboration has set limits for pair produced charge 2/3 or 4/3 third-generation leptoquarks decaying via $LQ \rightarrow \tau + b$ [5].

The data used for this analysis were collected by the D0 detector [6] operating at the Fermilab Tevatron Collider during the 1993–1996 period. The D0 detector is composed of three major systems: an inner detector for tracking charged particles, a uranium/liquid argon calorimeter for measuring electromagnetic and hadronic showers, and a muon spectrometer consisting of a magnetized iron toroid and three layers of drift tubes. The detector measures jets with an energy resolution of approximately $\sigma/E = 0.8/\sqrt{E}$ (E in GeV) and muons with a momentum resolution of $\sigma/p = [(\frac{0.18(p-2)}{p})^2 + (0.003p)^2]^{1/2}$ (p in GeV/ c). Missing transverse energy (\cancel{E}_T) is determined by summing the calorimeter and muon transverse energies, and is measured with a resolution of $\sigma = 1.08 \text{ GeV} + 0.019(\sum|E_{T,i}|)$ (E_T in GeV).

The decay of a b quark is indicated by the presence of a muon associated with a jet. We use three triggers to collect candidate leptoquark events, each requiring one or two muons [7]. A dimuon trigger required two muons with transverse momentum $p_T^\mu > 3.0$ GeV/ c . One single muon trigger required a muon with $p_T^\mu > 1.0$ GeV/ c and a jet with $E_T^j > 10$ GeV. The other single muon trigger, collected with a larger integrated luminosity, required a muon with $p_T^\mu > 10$ GeV/ c in the trigger and

$p_T^\mu > 15$ GeV/ c during off-line analysis, and a jet with $E_T^j > 15$ GeV. Integrated luminosities of 60.1, 19.5, and 92.4 pb $^{-1}$, respectively, were collected using these three triggers.

The off-line analysis uses muons in the pseudorapidity range $|\eta_\mu| < 1.0$ with $p_T^\mu > 3.5$ GeV/ c . The muon trajectories are required to be consistent with the reconstructed vertex position and have associated energy in the calorimeter. For events from either single muon trigger additional requirements are imposed: The presence of hits in all three muon detector layers, a matching track in the central detector, and a good fit [8] when these elements are combined. For the dimuon trigger events, at least one of the two muons must satisfy each of these additional requirements. Jets are reconstructed using only calorimeter energy with a cone algorithm of radius $R = 0.7$ where $R = \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2}$ about the jet's centroid and ϕ is the azimuthal angle. Each jet is required to have $E_T^j > 10$ GeV and to satisfy reconstruction quality criteria [9]. For the dimuon trigger, events with dimuon invariant mass greater than 8 GeV/ c^2 are selected to eliminate backgrounds from low mass resonances and each muon must be associated with a different jet with $\Delta R_{\mu\text{-jet}} < 0.5$ to increase b quark purity [10]. For the single muon triggers, we require a jet associated with the muon with the same ΔR requirement plus an additional jet with $E_T^j > 25$ GeV and $|n_j| < 1.5$.

We use \cancel{E}_T to identify neutrinos. To help eliminate events with poorly measured \cancel{E}_T , we reject events where the azimuthal angular separation, $\Delta\phi$, between the missing energy and the nearest jet is less than 0.7 rad. In some events, the source of the measured \cancel{E}_T is not mismeasured jets, but rather noise in individual calorimeter cells or from cells activated by the background generated by the Main Ring accelerator, which passes through the calorimeter. Such events are removed if the vector sum of the transverse energies in the jets and muons is consistent with zero.

Data from the two single muon triggers with large \cancel{E}_T have contributions from $W + \geq 2$ jet events where $W \rightarrow \mu\nu$ and the muon overlaps a jet. We use two variables, Z_μ and F_μ , for events passing those triggers to reduce the W boson acceptance. We define $Z_\mu \equiv p_T^\mu/H_T^{\mu,j}$, where $H_T^{\mu,j}$ is the scalar sum of the E_T of the jets and muons in the event. In Fig. 1 we compare the Z_μ distributions of data events that pass the low- p_T single muon trigger to Monte Carlo (MC) samples [11] satisfying a simulation of the trigger. There, for illustrative purposes, we select events with $\cancel{E}_T > 30$ GeV and $\Delta\phi > 0.6$ so that the data shown have roughly equal contributions from W boson and hadronic multijet events. Also shown are MC distributions of equal numbers of multijet events (which are b quark dominated due to their muon content) and $W \rightarrow \mu\nu$ events. The data (normalized to the total number of MC events) are consistent with the sum of these sources. Also shown is the same distribution




FIG. 1. The Z_μ distribution for events with $E_T > 30$ GeV and $\Delta\phi > 0.6$ from (a) data compared to MC events from multiple jet (solid) and $W \rightarrow \mu\nu$ (dashed) processes and (b) leptoquark events with $M_{LQ} = 100$ GeV/ c^2 . The arrow indicates the requirement $Z_\mu < 0.20$ used in this analysis.

for a leptoquark MC sample with $M_{LQ} = 100$ GeV/ c^2 satisfying the same criteria. Since the leptoquark Z_μ distribution is determined by b quark decay kinematics, it is similar to the multijet distribution. Requiring $Z_\mu < 0.20$ eliminates about 90% of the remaining $W \rightarrow \mu\nu$ background while maintaining a signal efficiency of 82% for $M_{LQ} = 100$ GeV/ c^2 . The second variable, F_μ , is the ratio of calorimeter energy within a cone of 0.4 centered on the muon's direction to that within a cone of 0.6: $F_\mu \equiv E(R_\mu \leq 0.4)/E(R_\mu \leq 0.6)$. Most hadronic energy in higher E_T direct b quark decays is spatially close to the muon. Requiring $F_\mu > 0.80$ removes about 84% of the $W \rightarrow \mu\nu$ background with a leptoquark signal efficiency of 82%. MC studies indicate that Z_μ and F_μ have little correlation.

The data from the high- p_T muon trigger include a significant contribution from top quark pair production ($t\bar{t}$) events. The scalar sum of jet E_T, H_T^j , was used to identify the top quark in Ref. [12]. In this analyses, we require $H_T^{\mu j} < 240$ GeV for those events satisfying the high- p_T single muon trigger to reduce the ($t\bar{t}$) contribution. Similarly, since the low- p_T single muon trigger has a larger contribution from multijet events, we reject events with six or more jets ($E_T^j > 10$ GeV) for that trigger.

The resulting E_T distribution for data from all three triggers after all selection criteria have been applied is given in Fig. 2. Also shown are the E_T distributions for MC events with $M_{LQ} = 100$ and 200 GeV/ c^2 . Requiring $E_T > 35$ GeV leaves two events.

We consider background contributions from $t\bar{t}$, intermediate vector boson, and multijet production of $b\bar{b}$ and $c\bar{c}$. Top quark events have multiple b quarks and E_T , but with additional, energetic jets. We use MC and our measured $t\bar{t}$ production cross section [13] to estimate that there are 1.4 ± 0.5 $t\bar{t}$ events in our sample. Intermediate vector boson events have E_T from $W \rightarrow l\nu$ or $Z \rightarrow \nu\bar{\nu}$ and muons near jets mimicking b quark decays when ei-




FIG. 2. The distributions of E_T for data events after all other selection requirements have been applied. Shown in the inset are E_T distributions for MC leptoquark events with leptoquark mass of 100 and 200 GeV/ c^2 .

ther a prompt muon overlaps a jet or a jet fragments into a muon via c quark or a π/K decay. Using our measured W and Z boson production cross sections [14] yields 1.0 ± 0.4 W boson events and 0.1 ± 0.1 Z boson events in this sample. Hadronically produced $b\bar{b}$ and $c\bar{c}$ events do not have energetic neutrinos and are effectively eliminated by the E_T and $\Delta\phi$ cuts. Estimates of their contribution using data and MC are consistent with zero, and we conservatively assume this in our limit calculation. Therefore, the total background is estimated to be 2.5 ± 0.6 events.

We calculate the detection efficiency for scalar leptoquark signals using MC acceptances multiplied by muon trigger and reconstruction efficiencies obtained from data samples collected using test triggers. The acceptances for different leptoquark masses are summarized in Table I. The use of a muon to tag b quark decays limits the acceptance to values under 3.5%. Factors contributing to this limited acceptance for the low- p_T single muon channel with $M_{LQ} = 100$ GeV/ c^2 include the muon branching

TABLE I. Third-generation scalar leptoquark acceptances for the three trigger channels and 95% C.L. limits on σB^2 for different M_{LQ} .

LQ mass (GeV/ c^2)	Acceptance ($\times 10^{-3}$)			σB^2 limit (pb)
	Dimuon	One muon low p_T	One muon High p_T	
50	0.04	0.95	0.14	144
60	0.08	2.4	0.35	59
80	0.29	6.0	0.93	22.6
100	0.40	9.7	1.9	12.7
125	0.59	14	2.8	8.9
150	1.2	18	3.2	7.0
200	1.3	24	3.1	6.0
250	1.9	27	3.7	5.1
300	1.8	31	2.2	5.4

fraction (0.35), muon and jet kinematic requirements (0.35), and muon trigger and reconstruction efficiency (0.25). The requirements used to reduce the background ($\cancel{E}_T, \Delta\phi, Z_\mu, F_\mu$) retain $\approx 40\%$ of the leptoquark signal.

We combine the three trigger channels to set limits. Errors on the acceptance are shown in Table II. Errors on trigger and reconstruction efficiencies are due to the statistical errors of the data used to calculate their values. Muon momentum resolution and jet energy scale errors are obtained from data, and their impact on the acceptance is determined using MC with $M_{LQ} = 100 \text{ GeV}/c^2$. The error on the F_μ cut efficiency is estimated by comparing data events without \cancel{E}_T requirements to MC multijet events. The three trigger channels have different systematic errors since their selection criteria and average muon p_T differ, but most errors are correlated. The total systematic error, including correlations and MC statistics, on the combined acceptance varies between 12.5% and 13.6% for different leptoquark masses.

The 95% confidence level (C.L.) upper limits on σB^2 include the systematic acceptance uncertainty and a 5.3% uncertainty in the integrated luminosity. The resulting upper limits for scalar leptoquark pair production as a function of leptoquark mass are given in Table I and shown in Fig. 3. Also shown in Fig. 3 are theoretical cross sections for the production of scalar and vector leptoquarks. The calculation of the scalar leptoquark cross section includes next-to-leading order diagrams and uses CTEQ4M parton distribution functions [15]. The theory band shown in the figure is determined using a renormalization factor of $\mu = M_{LQ}$ for the central value and $\mu = 2M_{LQ}$ and $M_{LQ}/2$ for the lower and upper bounds, respectively. The intersection of our limit curve with the lower edge of the theory based is at $94 \text{ GeV}/c^2$. This is our 95% C.L. lower limit on the mass of a charge-1/3 third-generation scalar leptoquark (taking $B = 1$ since $M_{LQ} < m_t$).

Similarly, we set limits for the mass of vector leptoquarks [16]. The vector leptoquark cross section has been calculated in the leading order approximation using CTEQ4M parton distribution functions and $\mu = M_{LQ}$ [17]. Vector leptoquarks are assumed to be either fundamental gauge bosons with Yang-Mills coupling or composite particles with anomalous coupling. For Yang-Mills

TABLE II. Summary of systematic errors in terms of % error on the acceptance.

Channel	Dimuon	One muon	
		low p_T	High p_T
Trigger	6.5	5.1	5.1
Reconstruction	5.1	5.7	4.2
Muon momentum resolution	1	1	10
Jet energy scale	4	4	2
F_μ cut	NA	10	6
$b \rightarrow \mu$ fraction	12	6	6




FIG. 3. The 95% C.L. limit on σB^2 (\bullet) compared to theoretical predictions. The prediction for scalar leptoquarks (solid line) use $\mu = 2M_{LQ}, M_{LQ}$, and $M_{LQ}/2$, while vector leptoquarks with minimal anomalous (dashed line) or Yang-Mills (dot-dashed line) coupling use $\mu = M_{LQ}$.

type coupling, a mass limit of $216 \text{ GeV}/c^2$ is obtained for $B = 1$. If $M_{LQ} > m_t$ the τt mode is allowed. We consider the case in which the branching fraction to $\nu_\tau b$ and τt each would be 0.5 if the fermion masses could be neglected relative to M_{LQ} . Taking into account phase space suppression factors [18], we determine that $M_{LQ} > 209 \text{ GeV}/c^2$ for Yang-Mills type vector leptoquarks for this $B < 1$ case. For anomalous coupling, we choose the coupling which yields the minimum pair production cross section. The intersection of our limit on σB^2 with the theory curve gives $M_{LQ} > 148 \text{ GeV}/c^2$ for minimal anomalous vector coupling.

In conclusion, we observe two events consistent with the final state $\nu\bar{\nu}b\bar{b}$ compared to an expected 2.5 ± 0.6 events from $t\bar{t}$ and W and Z boson production. We set limits on the mass of a charge-1/3 scalar or vector leptoquark. This result is independent of the coupling strength of a leptoquark to a third-generation lepton and quark.

We thank the staffs at Fermilab and collaborating institutions for their contributions to this work, and acknowledge support from the Department of Energy and National Science Foundation (U.S.A.), Commissariat à l'Énergie Atomique (France), State Committee for Science and Technology and Ministry for Atomic Energy (Russia), CAPES and CNPq (Brazil), Departments of Atomic Energy and Science and Education (India), Colciencias (Colombia), CONACyT (Mexico), Ministry of Education and KOSEF (Korea), and CONICET and UBA-CyT (Argentina).

*Visitor from Universidad San Francisco de Quito, Quito, Ecuador.

†Visitor from IHEP, Beijing, China.

- [1] J. L. Hewett and T. G. Rizzo, *Phys. Rep.* **183**, 193 (1989), and references therein.
- [2] M. Leurer, *Phys. Rev. D* **49**, 333 (1994).
- [3] B is defined as $1 - \beta$, where β is the leptoquark branching fraction into a charged lepton and quark.
- [4] OPAL Collaboration, G. Alexander *et al.*, *Phys. Lett. B* **263**, 123 (1991); L3 Collaboration, O. Adriani *et al.*, *Phys. Rep.* **236**, 1 (1991); ALEPH Collaboration, D. Decamp *et al.*, *Phys. Rep.* **216**, 253 (1992); DELPHI Collaboration, P. Abreu *et al.*, *Phys. Lett. B* **316**, 620 (1993).
- [5] CDF Collaboration, F. Abe *et al.*, *Phys. Rev. Lett.* **78**, 2906 (1997).
- [6] D0 Collaboration, S. Abachi *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **338**, 185 (1994).
- [7] M. A. C. Cummings, D. Hedin, and K. Johns, in *Proceedings of the Workshop on B Physics at Hadron Accelerators, Snowmass, CO, 1993*, edited by P. McBride and C. S. Mishra (SSCL, Dallas, TX, 1994).
- [8] T. Huehn, Ph.D. thesis, University of California-Riverside, 1995 (unpublished); see http://www-d0.fnal.gov/publications_talks/thesis/huehn/thesis.ps.
- [9] D0 Collaboration, S. Abachi *et al.*, *Phys. Rev. Lett.* **75**, 618 (1995).
- [10] D. Fein, Ph.D. thesis, University of Arizona, 1996 (unpublished); see http://www-d0.fnal.gov/publications_talks/thesis/fein/df_thesis.ps.
- [11] All Monte Carlo samples are generated with ISAJET and use a simulation of the detector, trigger, and off-line selections employing GEANT. F. Paige and S. Protopopescu, BNL Report No. BNL38034, 1986 (unpublished), release v6.49; R. Brun and F. Carminati, CERN Program Library Long Writeup W5013, 1993 (unpublished).
- [12] D0 Collaboration, S. Abachi *et al.*, *Phys. Rev. Lett.* **74**, 2632 (1995).
- [13] D0 Collaboration, S. Abachi *et al.*, *Phys. Rev. Lett.* **79**, 1203 (1997). We use a top quark mass of $170 \text{ GeV}/c^2$ for background calculations.
- [14] D0 Collaboration, S. Abachi *et al.*, *Phys. Rev. Lett.* **75**, 1456 (1995).
- [15] M. Krämer, T. Plehn, and P. M. Zerwas, *Phys. Rev. Lett.* **79**, 341 (1997).
- [16] For masses greater than $150 \text{ GeV}/c^2$, MC acceptances for vector leptoquarks are found to be greater than or equal to those for scalar leptoquarks with the same M_{LQ} .
- [17] J. Blumlein, E. Boos, and A. Kryukov, *Z. Phys. C* **76**, 137 (1997). Leading order calculations with $\mu = M_{LQ}$ yield about the same cross section as next-to-leading order with $\mu = 2M_{LQ}$.
- [18] We use $\sqrt{(1 + d_1 - d_2)^2 - 4d_1} [1 - (d_1 + d_2)/2 - (d_1 - d_2)^2/2]$ with $d_1 = (m_t/M_{LQ})^2$ and $d_2 = (m_\tau/M_{LQ})^2$. T. Rizzo (private communication).